

Weekly Assessment for Delta Operations on ESA and CESA-listed Salmonids and Osmerids including Current Delta Hydrologic Conditions

Final: Tuesday, January 13, 2026 at 4 PM

Table of contents

Executive Summary	2
ESA and CESA-listed Salmonids	2
ESA and CESA-listed Osmerids	2
1 Current Delta Hydrologic Conditions	2
1.1 Operational and Regulatory Conditions	2
1.2 Current Conditions	2
1.3 Zone of Influence	4
2 Assessment for Delta Operations on Salmonids	5
2.1 Natural Winter-run Chinook	5
2.1.1 Juvenile Production Estimate	5
2.1.2 Current Status	5
2.1.3 Annual Loss	6
2.1.4 STARS	6
2.2 Hatchery Winter-run Chinook	8
2.2.1 Hatchery Releases	8
2.2.2 Juvenile Production Estimate	8
2.2.3 Annual Loss	8
2.3 Natural-origin Central Valley Steelhead	9
2.3.1 Current Status	9
2.3.2 Annual Loss	9
2.4 Hatchery-origin Central Valley Steelhead	10
2.4.1 Surrogate Releases	10

2.5	Spring-run	11
2.5.1	Current Status	11
2.5.2	Spring-run Surrogate Releases	12
2.5.3	Annual Loss	12
2.6	Loss Prediction and Trajectories	13
2.7	Evaluation	13
3	Weekly Assessment for Delta Operations on ESA and CESA-listed Osmerids	13
3.1	Operational and Regulatory Conditions	13
3.2	Delta smelt	13
3.2.1	Biological	13
3.2.2	Environmental	17
3.2.3	Real-time Assessment Thresholds	17
3.2.4	Evaluation	19
3.3	Longfin smelt	19
3.3.1	Biological	19
3.3.2	Real-time Assessment Thresholds	23
3.3.3	Evaluation	23
3.4	End of smelt Entrainment Management	24
3.5	References	24

Executive Summary

ESA and CESA-listed Salmonids

- Entrainment management season is **active**.
- Season Loss: **0** (0.00% of threshold) DNA Winter-run, **0** (0.00% of threshold) Hatchery Winter-run, **43.91** (0.83% of threshold) Natural Steelhead, **122.07** (3.42% of threshold) Hatchery Steelhead, and **872.69** (10.84% of threshold) Spring-run Surrogates.
- Winter-run presence in the Delta is **high** (historical peak).
- Steelhead presence in the Delta is **increasing**.

ESA and CESA-listed Osmerids

- The First flush action implementation period ended on 1/7/26. Entrainment management is currently active.
- Delta smelt are primarily distributed west of the confluence, in Suisun Marsh
- No Delta smelt or longfin smelt salvage has been observed this water year
- Turbidity in the central/south Delta is moderate.

1 Current Delta Hydrologic Conditions

1.1 Operational and Regulatory Conditions

Entrainment management is the current controlling factor. See most recent weekly outlook for more information.

1.2 Current Conditions

Most recent inflow at Freeport in the Sacramento River and Vernalis in the San Joaquin River is 65,994 and 2,951 cfs respectively. Most recent 1-day, 5-day, and 14-day OMRI measurements were -4,900, -4,909, and -5,061 respectively, and most recent export data were 3,551 for Jones Pumping Plant and 1,734 for Henry O. Banks Pumping Plant.




Figure 1: Operations and Action Summary, WY 2026. The numbers and dashed lines in the OMRI plot indicate different triggers (see Table 1), with shading representing specific action periods. OMRI data (colored lines) calculated by SacPAS, Freeport (FPT) and Vernalis (VNS) flow data from CDEC, and CVP (TRP) and SWP (HRO) exports data from CDEC.

Table 1: Summary of Actions and Triggers, WY 2026

La- bel	Action	Date Triggered	Date Implemented	Number Days Implemented	Regulation
1	DCC Gate Closure	10/28/2025	2025-10-30	Ongoing	DCC gates
2	First Flush	12/24/2025	2025-12-25	14 days	Entrainment Management

1.3 Zone of Influence

Zone of Influence (ZOI) analysis is discussed in detail in the December 22 assessment. Current conditions were queried from most recent Freeport flow data on the Sacramento River and Vernalis flow data on the San Joaquin river from [SacPAS](#). Forecasted flows were queried from short range deterministic flows provided by the [California Nevada River Forecast Center](#).

Current conditions at Freeport and Vernalis indicate that delta hydrology falls within the 'himed' category. Forecasted conditions averaged across the next 7 days falls within the 'himed' category.

The altered channel length for the current "himed" hydrology is 23, 53, 118 and 111 kilometers (km) across OMR bins of -2000, -3500, -5000 and <-5500 respectively. The altered channel length for forecasted "himed" hydrology is 23, 53, 118 and 111 kilometers (km) across OMR bins of -2000, -3500, -5000 and <-5500 respectively.

Change in altered channel length between OMR levels is 88 km for current conditions and 88 km for forecasted conditions indicating that ZOI impacts across OMR scenarios would not change between current and forecasted conditions. Across the nine hydrology bins, changes in altered channel length across OMR scenarios are moderate (between 25th and 75th percentiles) for both current and forecasted hydrology.




Figure 2: Modeled Zone of Influence at different OMR scenarios based on current inflow hydrology (left) and forecasted inflow hydrology (right) from the Sacramento River and San Joaquin River

2 Assessment for Delta Operations on Salmonids

For more detailed data on salmonid conditions in the Delta see corresponding webpage on [SacPAS](#).

2.1 Natural Winter-run Chinook

2.1.1 Juvenile Production Estimate

The Juvenile Production Estimate for winter-run is 1,057,452 for the current water year.

2.1.2 Current Status

Entry Timing - Historically, as of Jan 12, 75% of length-at-date (LAD) winter-run have entered the delta based on Knights Landing RST catch, 2% have exited the delta based on Chipps Island Trawl Catch, and 0% of DNA confirmed winter-run have been salvaged.

Table 2

Species	Red Bluff Diversion Dam	Tis- dale RST	Knights Landing RST	Sac Trawl (Sher- wood)	Chipps Island Trawl	Sal- vage
Chinook, LAD	98%	78%	75%	38%	2%	15%
Winter-run, Unclipped						
Chinook, DNA	NA	NA	NA	NA	NA	0%
Winter-run, Unclipped (Water Year)						

Red Bluff Diversion Dam Passage Estimate - As of Dec 16 estimated passage to date of LAD winter run at Red Bluff Diversion is approximately 3.61 million fish. * *Note that outmigration timing overlaps with spring run migrating fish, and true winter-run abundance likely differs from these estimates.*

Delta Monitoring - Total catch of LAD winter run at RSTs at Delta Entry (Tisdale, Knights Landing, Lower Sacramento River) between Dec 30 and Jan 12 is 190 individuals. Total catch at Sacramento Trawl and Beach Seines in the delta between Dec 30 and Jan 12 is 15 individuals. Total catch at Delta Exit at Chipps Island between Dec 30 and Jan 08 is 1 individuals.

2.1.3 Annual Loss

The annual Loss threshold for natural winter-run is 1% of the jpe or 10,574.52 fish. As of January 12, cumulative loss of genetically confirmed winter-run is 0 or 0.00% of the annual loss threshold. Cumulative loss in the past 7 days has been 0.




Figure 3: Cumulative loss of natural-origin winter-run for WY 2026. Cumulative loss is based on genetically confirmed winter-run captured in salvage or length-at-date winter-run in which genetic confirmation was unable to be obtained

2.1.4 STARS

The Delta STARS Model is an individual-based simulation model that predicts survival, travel time, and routing of juvenile salmon migrating through the Sacramento–San Joaquin River Delta. This model gives insight into survival and routing patterns of winter-run based on most current conditions.

As of January 12, overall through delta STARS estimated survival probability (with 80% credible intervals) is 0.73 (0.66-0.8) placing it in the 94th percentile of historical STARS survival estimates for the month of January (WYs 2018-2025). STARS estimated routing and survival probabilities (with 80% credible intervals) into the interior delta are 0.09 (0.08-0.11) and 0.61 (0.45-0.76), respectively, corresponding to the 6th and 90th percentiles of historical January estimates (WYs 2018-2025).




Figure 4: Estimated overall winter-run survival from Knights Landing to Chipps Island. Black line indicates the current water-year, and other colored lines correspond to past water years.




Figure 5: Estimated survival from Knights Landing to Chipps Island of simulate winter-run cohorts that route through the interior delta. Black line indicates the current water-year, and other colored lines correspond to past water years.




Figure 6: Estimated probability of winter-run routing into the interior delta. Black line indicates the current water-year, and other colored lines correspond to past water years.

2.2 Hatchery Winter-run Chinook

2.2.1 Hatchery Releases

To date, no winter-run Livingstone hatchery releases have occurred in WY 2026

2.2.2 Juvenile Production Estimate

The Juvenile Production Estimate for hatchery winter-run is 130,096 for Livingston Stone releases.

2.2.3 Annual Loss

To date, no loss has occurred as no hatchery winter-run have been released.

2.3 Natural-origin Central Valley Steelhead

2.3.1 Current Status

Delta Entry Timing - Historically, as of Jan 12, 16% of CCV steelhead have entered the delta based on Knights Landing RST catch, 1% have exited the delta based on Chipps Island Trawl Catch, and 6% have been salvaged.


Table 3: Average Percent of annual emigrating population for unclipped CCV steelhead captured at the following locations and salvaged at SWP and CVP Delta facilities for the past 10 years.

Species	Red Bluff Diversion Dam	Tisdale RST	Knights Landing RST	Sac Trawl (Sherwood)	Chipps Island Trawl	Sal- vage
Steelhead, Unclipped		1%	15%	16%	0%	1% 6%

Delta Monitoring - Total catch of LAD winter run at RSTs at Delta Entry (Tisdale, Knights Landing, Lower Sacramento River) between Dec 30 and Jan 12 is 0 individuals. Total catch at Sacramento Trawl and Beach Seines in the delta between Dec 30 and Jan 12 is 0 individuals. Total catch at Delta Exit at Chipps Island between Dec 30 and Jan 08 is 0 individuals.

2.3.2 Annual Loss

As of January 12, cumulative loss of unclipped steelhead is 43.91 or 0.83% of the incidental take limit in the NMFS Biological Opinion. Cumulative loss in the past 7 days has been 97.59.



www.cbr.washington.edu/sacramento/

13 Jan 2026 15:49:02 PST

Figure 7: Cumulative loss of natural-origin steelhead for WY 2026.

2.4 Hatchery-origin Central Valley Steelhead

2.4.1 Surrogate Releases

There have been a total of 2 releases totaling 878,848 steelhead in Water Year 2026. JPE for the hatchery releases as of today is 356,526 based on estimated survivals using forecasted water year types (see details in table below). The annual loss threshold, equal to 1% of the JPE, is currently 3565, but is subject to change with additional steelhead releases.

Table 4: Summary of steelhead hatchery releases in Water Year 2026

Hatchery	Date of Release	Mean Fork Length (mm)	Number Released	Estimated Survival	Juvenile Production Estimate
Nimbus	2025-11-02	223	233,109	72%	167,838
Coleman	2025-12-19	195	645,739	29%	188,688

Total loss of hatchery-origin steelhead is 122.07 or 3.42% of the threshold. *Note that hatchery origin of salvaged fish can not be determined at this time and salvage is based on the assumption of similar routing and survival probabilities of individual hatchery releases.




Figure 8: Cumulative loss of hatchery steelhead for WY 2026.

2.5 Spring-run

2.5.1 Current Status

Delta Entry Timing- Historically, as of Jan 12, 26% of LAD spring run have entered the delta based on Knights Landing RST catch, 0% have exited the delta based on Chipps Island Trawl Catch, and 0% have been salvaged.

Table 5: Average Percent of annual emigrating population for LAD Spring-run Chinook Salmon captured at the following locations and salvaged at SWP and CVP Delta facilities for the past 10 years.

Species	Red Bluff Diversion Dam	Tis- dale RST	Knights Landing RST	Sac Trawl (Sherwood)	Chipp Island Trawl	Sal- vage
Chinook, LAD Spring-run, Unclipped	12%	14%	26%	2%	0%	0%

Red Bluff Diversion Dam Passage Estimate - As of Dec 16 estimated passage to date of LAD spring run at Red Bluff Diversion is approximately 0.01 million fish. * Note that outmigration

timing overlaps with winter run and fall run outmigration, and true spring run abundance likely differs from these estimates.

Delta Monitoring- Total catch of LAD winter run at RSTs at Delta Entry (Tisdale, Knights Landing, Lower Sacramento River) between Dec 30 and Jan 12 is 172 individuals. Total catch at Sacramento Trawl and Beach Seines in the delta between Dec 30 and Jan 12 is 0 individuals. Total catch at Delta Exit at Chipps Island between Dec 30 and Jan 08 is 0 individuals.

2.5.2 Spring-run Surrogate Releases

A total of 805,323 spring-run surrogate fish have been released in Water Year 2026. See details in table below.

Table 6: Spring-run Chinook Salmon Surrogate Releases.

Hatchery	Release Date	Type	# of CWT Fish Released	Confirmed Loss	CWT Codes
Cole- man NFH	2025-11- 13	Year- ling	143346	8.7	056808, 056809
Cole- man NFH	2025-11- 17	Year- ling	75119	0.0	056810
Cole- man NFH	2025-12- 17	Year- ling	468876	639.9	053700, 056806, 056811, 056812, 056814, 056815, 056817
Cole- man NFH	2025-12- 22	Year- ling	60873	224.1	056813
Cole- man NFH	2026-01- 08	Year- ling	57109	0.0	056816

2.5.3 Annual Loss

The annual loss threshold is 1% of the total releases, which equals 8,053 fish. As of January 12, cumulative loss is 872.69 fish or 10.84% of the annual loss threshold.

2.6 Loss Prediction and Trajectories

The following figures display the cumulative loss trajectories relative to historical years (“Spaghetti Plots”) and the current loss predictor model outputs for Winter-run Chinook Salmon and Steelhead.

2.7 Evaluation

1. **What is the probability of exceeding natural or hatchery winter-run Chinook Salmon loss thresholds in the upcoming week?**

LOW RISK: Cumulative loss is currently 0% of the threshold. Current trajectory suggests the threshold is unlikely to be exceeded in the upcoming week.

2. **What is the probability of spring-run hatchery Chinook Salmon loss thresholds in the upcoming week?**

LOW RISK: Cumulative loss is currently 10.8% of the threshold. Current trajectory suggests the threshold is unlikely to be exceeded in the upcoming week.

3. **What is the probability of hatchery Steelhead loss thresholds in the upcoming week?**

LOW RISK: Cumulative loss is currently 3.4% of the threshold. Current trajectory suggests the threshold is unlikely to be exceeded in the upcoming week.

3 Weekly Assessment for Delta Operations on ESA and CESA-listed Osmerids

3.1 Operational and Regulatory Conditions

- See current Weekly Fish and Water Operations Outlook document.
- Additional information also available on the [SacPAS SMT page](#).

3.2 Delta smelt

3.2.1 Biological

- **Delta smelt life stages:** Adult, Juvenile
- **Abundance estimate:** 8963 (95% CL: 1,984 to 26,236) as of the week of January 5–9, 2026




Figure 9: Natural LAD Winter-Run Chinook Salmon Loss Cumulative to Date with historical years and Single Year Loss Thresholds.




Figure 10: Natural Central Valley Steelhead Loss Cumulative to Date with historical years and Single Year Loss Thresholds.




Figure 11: Estimates of Winter-run Chinook Loss generated by Loss and Salvage Predictor tool.




Figure 12: Estimates of Steelhead Loss generated by Loss and Salvage Predictor tool.

- **Releases:** A total of 163,349 cultured Delta smelt have been released for WY 2026. The most recent release of 24,606 fish occurred in Sacramento River at Rio Vista on Dec 16, 2025.
- **Delta smelt count:** 33 adult Delta smelt and 24 juvenile Delta smelt have been detected this water year. See Table 7 for recent detections, Figure 13 for spatial distribution, and Figure 14 for temporal distribution.
- **Delta smelt salvage:** 0 Delta smelt have been salvaged, and the cumulative seasonal salvage is 0.

Notes

- Since there are few recent detections of Delta smelt, estimation of distribution within the Delta is limited.
- As mentioned in EDSM reporting, fork length ranges reported for Delta smelt and longfin smelt life stages are defined by permit reporting purposes and are not intended to delineate cohorts or distinguish from hatchery or wild origin. See Table 7 caption for fork-length ranges for age groups of Delta smelt.
- See [SacPAS SMT Page](#) for additional details on releases and detection in surveys and salvage.
- Historical salvage trends can be found at: [SacPAS Salvage Timing](#)

Table 7: Delta smelt detections in the last 2 weeks. Fork Length > 58mm = Adult, Fork Length 20-58mm = Juvenile, Fork Length < 20mm = Larva.

Survey	Date	Region	Stratum	Life Stage	Catch
EDSM	2026-01-02	West	Suisun Marsh	Adult	2
EDSM	2026-01-02	West	Suisun Marsh	Juvenile	1
EDSM	2026-01-07	West	Suisun Marsh	Adult	1
EDSM	2026-01-07	West	Suisun Marsh	Juvenile	1
EDSM	2026-01-08	West	Suisun Marsh	Adult	3
EDSM	2026-01-08	West	Suisun Marsh	Juvenile	1





Figure 13: Delta smelt distribution for WY 2026

Table 8: Delta smelt water year totals by life stage

Survey	Region	Life Stage	Total
DJFMP	North	Juvenile	1

Table 8: Delta smelt water year totals by life stage

Survey	Region	Life Stage	Total
EDSM	North	Adult	1
EDSM	West	Adult	32
EDSM	West	Juvenile	23




Figure 14: Time series of Delta smelt catch, WY 2026

3.2.2 Environmental

3.2.2.1 First Flush

- Not relevant

3.2.3 Real-time Assessment Thresholds

3.2.3.1 Adult Delta smelt


Threshold: If daily average JPF < 0 AND turbidity > 12 FNU at OBI, HOL and OSJ

- JPF: 9,041 cfs as of Jan 12, 2026

- **OBI Turbidity:** 6.76, 6.66, 6.46 FNU as of Jan 12, 2026
- **HOL Turbidity:** 8.52, 7.39, 7.01 FNU as of Jan 12, 2026
- **OSJ Turbidity:** 15.31, 18.22, 18.8 FNU as of Jan 12, 2026

Offramp Adult Protections when RVB or SJJ > 12°C

- **RVB temperature (3-day average):** 9.98°C as of Jan 12, 2026



Preliminary data from CDEC; subject to revision.

www.cbr.washington.edu/sacramento/
13 Jan 2026 12:35:02 PST

- See [Bay-Delta Live](#) for recent Delta-wide turbidity conditions.

3.2.3.2 Larval/juvenile Delta smelt

Threshold: After the onset of spawning, if $JPF < 0$ cfs AND turbidity is ≥ 12 FNU in the south Delta AND PTM modeling indicates the action would avoid $\geq 5\%$ entrainment of Delta smelt population after 30 days

- **12-station South Delta Turbidity:** The most recent average turbidity was 19.1 FNU as of Dec 30, 2025

3.2.4 Evaluation

Delta smelt:

1. After the start of entrainment management, is $JPF < 0$ and is daily average turbidity ≥ 12 FNU in the OMR corridor (stations OBI, HOL, and OSJ)?

No, these conditions will not be met this week.

2. Has the average water temperature at Jersey Point or Rio Vista not exceeded 53.6°F (12°C) for 3 consecutive days and/or has this action already been taken during WY 2026?

Temperature at Rio Vista or Jersey Point has not exceeded the threshold. The Delta smelt adult entrainment management action has not yet been taken in WY 2026.

3. What is the evidence for the onset of Delta smelt spawning?

Upstream migration for Delta smelt occurs between September and December and in response to “first flush” conditions (Sommer et al. 2011, Grimaldo et al. 2009). Migration typically ranges one to four weeks after flow and turbidity increases, based on salvage data (Sommer et al. 2011). Historically, detections of ripe Delta smelt began in January and peaked in February and March and the majority of Delta Smelt spawning occurs within a temperature range of $9\text{--}18^{\circ}\text{C}$ (Damon et al. 2016). Based on [historical monitoring data](#) from the past few years, first detection of larvae in the Central and South Delta has typically occurred by mid to late March. Because first flush conditions were met on December 23, spawning-related behaviors may begin occurring during the current assessment week, consistent with the typical one- to four-week response window following increased flow and turbidity.

4. After the onset of spawning, have the following conditions occurred: $\geq 5\%$ entrainment of the Delta smelt population at facilities after 30 days?

Although spawning may begin during the current assessment period, JPF is above 0 cfs; therefore, the conditions required to evaluate larval and juvenile Delta smelt entrainment management are not met.

3.3 Longfin smelt

3.3.1 Biological

- **Longfin smelt life stages:** Juvenile, Adult
- **Longfin smelt count:** 296 adult, 446 juvenile, and 10 larval longfin smelt have been detected this water year. See Table 9 for recent detections, Figure 15 for spatial distribution, and Figure 16 for temporal distribution.

- **Longfin smelt salvage:** 0 longfin smelt have been salvaged, and the cumulative seasonal salvage is 0.

Table 9: Longfin smelt detections in the last 2 weeks. Fork Length > 84mm = Adult, Fork Length 20-84mm = Juvenile, Fork Length < 20mm = Larva.

Survey	Date	Region	Stratum	Life Stage	Catch
DJFMP	2025-12-31	N/A	Chippis Island	Adult	44
DJFMP	2025-12-31	N/A	Chippis Island	Juvenile	2
DJFMP	2026-01-05	Bay	Bay	Juvenile	1
DJFMP	2026-01-05	N/A	Chippis Island	Adult	1
DJFMP	2026-01-06	N/A	Chippis Island	Adult	1
DJFMP	2026-01-08	N/A	Chippis Island	Adult	3
EDSM	2025-12-31	Far West	Suisun Bay	Adult	2
EDSM	2025-12-31	Far West	Suisun Bay	Juvenile	1
EDSM	2026-01-05	Far West	Suisun Bay	Adult	2
EDSM	2026-01-05	Far West	Suisun Bay	Juvenile	2
EDSM	2026-01-06	West	Suisun Bay	Juvenile	1
EDSM	2026-01-07	West	Suisun Marsh	Juvenile	1
EDSM	2026-01-12	North	Lower Sacramento	Juvenile	1
EDSM	2026-01-12	West	Lower Sacramento	Adult	19
EDSM	2026-01-12	West	Lower Sacramento	Juvenile	5





Figure 15: Longfin Smelt Distribution for WY 2026

Table 10: Longfin smelt water year totals by life stage

Survey	Region	Life Stage	Total
DJFMP	Bay	Juvenile	1

Table 10: Longfin smelt water year totals by life stage

Survey	Region	Life Stage	Total
DJFMP	N/A	Adult	240
DJFMP	N/A	Juvenile	14
EDSM	Far West	Adult	7
EDSM	Far West	Juvenile	13
EDSM	North	Juvenile	1
EDSM	West	Adult	41
EDSM	West	Juvenile	80
baystudy	Bay	Adult	6
baystudy	Bay	Juvenile	320
baystudy	Far West	Adult	2
baystudy	Far West	Juvenile	11
baystudy	West	Juvenile	6
sls	Far West	Larva	3
sls	North	Larva	1
sls	West	Larva	6




Figure 16: Time series of longfin smelt catch, WY 2026

3.3.2 Real-time Assessment Thresholds

3.3.2.1 Start of Entrainment Management (Adult Longfin Smelt)

- Not relevant

3.3.2.2 Adult longfin smelt

- **Threshold:** $JPF < 0$ cfs, annual loss is on a trajectory to exceed 5% of the adult population abundance, and reduced exports will reduce entrainment in the south Delta
 - Daily average $JPF: 9,041$ cfs as of Jan 12, 2026
 - Adult abundance (Age 1+ LFS index): 2479.2 fish
 - * 5% of abundance + 1: 125.0
 - Water year total adult longfin smelt salvage = 0

3.3.2.3 Larval/juvenile longfin smelt

- **Threshold:** $JPF < 0$ cfs AND population model demonstrates need to reduce entrainment to avoid population decline
 - Daily average $JPF: 9,041$ cfs as of Jan 12, 2026

3.3.3 Evaluation

Longfin smelt:

1. If $JPF < 0$, what is the trajectory of annual loss of adult longfin smelt and is it likely to exceed 5% of the adult population estimate? Is South Delta entrainment expected to decrease due to a reduction in export pumping?

JPF is not < 0 cfs and no adult longfin smelt have been detected in salvage.

2. For larval and juvenile longfin smelt, if $JPF < 0$ cfs, do particle tracking models show a moderate to high difference in particle fates across different OMRI scenarios? Does Zone of Influence modeling show moderate to high changes in hydrodynamic footprint across different OMRI scenarios? Are these effects anticipated to cause a population decline?

JPF is not less than zero. ZOI modeling shows moderate change in the hydrodynamic footprint between OMRI scenarios.

3. Is there additional information or other analyses that should be considered in this evaluation?

Additional information may be discussed if needed at the DAT call.

3.4 End of smelt Entrainment Management

- Not relevant

3.5 References

Damon, L. J., S. B. Slater, R. D. Baxter, and R. W. Fujimura. 2016. Fecundity and reproductive potential of wild female Delta smelt in the upper San Francisco Estuary, California. California Fish and Game 102(4):188–210.

Grimaldo, L. F., T. Sommer, N. Van Ark, G. Jones, E. Holland, P. B. Moyle, B. Herbold & P. Smith (2009) Factors Affecting Fish Entrainment into Massive Water Diversions in a Tidal Freshwater Estuary: Can Fish Losses be Managed? North American Journal of Fisheries Management, 29:5, 1253-1270, DOI: 10.1577/M08-062.1

Sommer, T., F. Mejia, M. Nobriga, and L. Grimaldo. 2011. The Spawning Migration of Delta Smelt in the Upper San Francisco Estuary. San Francisco Estuary and Watershed Science 9(2).